

Basis and Dimension

Recall Example 10 (a) from the last set of notes. In this example, we saw a collection of two vectors that were a spanning set for  $\mathbb{R}^2$ . We noted that if we added vectors to this set, it would still form a spanning set. However, this would create sort of “trivial” spanning sets. What we desire is a “minimal” spanning set for a vector space. This is our goal in this set of notes. Before we get to this, we need a definition.

**Example 1:** Consider the set of vectors  $\{x_1, x_2, x_3\}$  in  $\mathbb{R}^3$  where  $x_1 = \begin{bmatrix} 2 \\ -1 \\ 5 \end{bmatrix}$ ,

$x_2 = \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix}$ , and  $x_3 = \begin{bmatrix} -1 \\ 11 \\ -7 \end{bmatrix}$ . Let  $S = \text{Span}(x_1, x_2, x_3)$ . If we pick just two of

these vectors, say  $x_1$  and  $x_2$ , it is clear that  $\text{Span}(x_1, x_2) \subseteq S$ . In fact,  $\text{Span}(x_1, x_2)$  is a *subspace* of  $S$ . Notice that  $x_3$  is a linear combination of the other two vectors. In particular,

$$x_3 = -2x_1 + 3x_2. \tag{1}$$

Therefore, every vector that is a linear combination of  $\{x_1, x_2, x_3\}$  can actually be written as a linear combination of just  $x_1$  and  $x_2$ :

$$\begin{aligned} v &= a_1x_1 + a_2x_2 + a_3x_3 \\ &= a_1x_1 + a_2x_2 + a_3(-2x_1 + 3x_2) \\ &= (a_1 - 2a_3)x_1 + (a_2 + 3a_3)x_2 \end{aligned}$$

What this means is that  $\text{Span}(x_1, x_2) = \text{Span}(x_1, x_2, x_3)$ . But we could rewrite Equation (1) as

$$2x_1 - 3x_2 + x_3 = 0 \tag{2}$$

and see that we could solve for any one of the three vectors (since all coefficients are non-zero). It follows that

$$\text{Span}(x_1, x_2, x_3) = \text{Span}(x_1, x_2) = \text{Span}(x_2, x_3) = \text{Span}(x_1, x_3).$$

This is because of the dependency relationship represented by Equation (2).

On the other hand, no such relationship exists between  $x_1$  and  $x_2$ . If there was one, there would exist coefficients  $c_1$  and  $c_2$ , not both zero, such that

$$c_1x_1 + c_2x_2 = 0. \quad (3)$$

We could then solve for one of the vectors in terms of the other and see that the two vectors were multiples of one another (which isn't the case). Therefore, the only way Equation (3) can hold is if  $c_1 = c_2 = 0$ . As is usually the case in mathematics, this important characteristic has a name.

**Definition:** Let  $V$  be a vector space. The vectors  $v_1, v_2, \dots, v_n$  are said to be **linearly independent** if the only solution to

$$c_1v_1 + c_2v_2 + \dots + c_nv_n = 0$$

is  $c_1 = c_2 = \dots = c_n = 0$ . If there is a solution in which not *all* coefficients are zero, then the vectors are **linearly dependent**.

**Definition:** Let  $V$  be a vector space. The vectors  $v_1, v_2, \dots, v_n$  form a **basis** of  $V$  if:

- (a)  $v_1, v_2, \dots, v_n$  are linearly independent, and
- (b)  $v_1, v_2, \dots, v_n$  span  $V$ .

**Example 2:** Are the vectors  $\begin{bmatrix} 2 \\ 1 \end{bmatrix}$  and  $\begin{bmatrix} 3 \\ 2 \end{bmatrix}$  linearly independent or dependent?

Consider  $c_1 \begin{bmatrix} 2 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} 3 \\ 2 \end{bmatrix} = 0$ . This implies that  $2c_1 + 3c_2 = 0$  and  $c_1 + 2c_2 = 0$ .

Solving this system of equations, we see that  $c_1 = c_2 = 0$  is the only solution. So they are independent.

**Example 3:** Are the vectors  $\begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix}$ ,  $\begin{bmatrix} 3 \\ 0 \\ 2 \end{bmatrix}$  and  $\begin{bmatrix} 10 \\ -2 \\ 12 \end{bmatrix}$  linearly independent or dependent?

As before, if  $c_1 \begin{bmatrix} 2 \\ -1 \\ 4 \end{bmatrix} + c_2 \begin{bmatrix} 3 \\ 0 \\ 2 \end{bmatrix} + c_3 \begin{bmatrix} 10 \\ -2 \\ 12 \end{bmatrix} = 0$ , then we have the system of equations

$$\begin{aligned} 2c_1 + 3c_2 + 10c_3 &= 0 \\ -c_1 - 2c_3 &= 0 \\ 4c_1 + 2c_2 + 12c_3 &= 0 \end{aligned}$$

Solving this, we have solution set  $\{(-2\alpha, -2\alpha, \alpha) : \alpha \in \mathbb{R}\}$ . So they are dependent.

As you may have noticed, determining whether or not a set of vectors is linearly independent or linearly dependent boils down to solving a system of equations. We have already seen how this is related to the determinant of a matrix.

**Theorem 1:** Let  $v_1, v_2, \dots, v_n$  be vectors in  $\mathbb{R}^n$  and let  $A$  be the  $n \times n$  matrix whose  $i^{\text{th}}$  column is  $v_i$ . Then  $v_1, v_2, \dots, v_n$  are linearly dependent if and only if  $A$  is singular.

**Theorem 2:** Let  $V$  be a vector space with  $\{v_1, v_2, \dots, v_n\}$  a spanning set. Then any collection of  $m$  vectors, where  $m > n$ , is linearly dependent.

**Corollary:** Let  $V$  be a vector space and suppose that  $\{v_1, v_2, \dots, v_n\}$  and  $\{u_1, u_2, \dots, u_m\}$  are both bases for  $V$ . Then  $n = m$ .

This last corollary tells us that the number of vectors in a basis is a fixed number for each vector space. Let's name it.

**Definition:** Let  $V$  be a vector space. The number of vectors in a basis is called the *dimension* of  $V$  (this can be finite or infinite). We say the trivial vector space  $\{0\}$  has dimension 0.

**Example 4:**  $\mathbb{R}^3$  is a finite-dimensional vector space of dimension 3. The vector space consisting of all polynomials, usually denoted by  $P$ , is an infinite dimensional vector space.

**Theorem 3:** Let  $V$  be a vector space of dimension  $n > 0$ . Then:

- (a) Any set of  $n$  linearly independent vectors spans  $V$ .
- (b) Any set of  $n$  vectors that span  $V$  are linearly independent.

**Proof:** (a) Let  $v_1, v_2, \dots, v_n$  be linearly independent and let  $v$  be any vector of  $V$ . By Theorem 2, the vectors  $v_1, v_2, \dots, v_n, v$  are linearly dependent. So there exist scalars  $c_1, c_2, \dots, c_n, c_{n+1}$ , not all zero, for which  $c_1 v_1 + c_2 v_2 + \dots + c_n v_n + c_{n+1} v = 0$ . If  $c_{n+1} = 0$ , then this gives us a nontrivial linear combination of  $v_1, v_2, \dots, v_n$  that is the zero vector which would imply that  $v_1, v_2, \dots, v_n$  are linearly dependent. So  $c_{n+1} \neq 0$ , and therefore

$$v = \left(-\frac{c_1}{c_{n+1}}\right)v_1 + \left(-\frac{c_2}{c_{n+1}}\right)v_2 + \dots + \left(-\frac{c_n}{c_{n+1}}\right)v_n.$$

Hence,  $v \in \text{Span}(v_1, v_2, \dots, v_n)$ . Since  $v$  was arbitrary, we see that  $v_1, v_2, \dots, v_n$  span  $V$ .

(b) Suppose  $v_1, v_2, \dots, v_n$  span  $V$  and suppose that they are linearly dependent. Then one of the vectors, say  $v_n$ , can be written as a linear combination of the others. This means that  $v_1, v_2, \dots, v_{n-1}$  also span  $V$ . If these vectors are also linearly dependent, we can eliminate another vector. Eventually, we will arrive at a set of  $k$  vectors ( $k < n$ ) that (1) still span  $V$ , and (2) are finally linearly independent. But this contradicts the corollary Theorem 2. So the vectors  $v_1, v_2, \dots, v_n$  must already be linearly independent.

Let's notice the impact of this theorem. The definition of "basis" says that two conditions must be met for a set of vectors to be a basis: they must be linearly independent and they must span the vector space. However, this theorem tells us that if we have the right number of vectors, either condition is enough. In fact, in the proof, we saw even more than that.

**Theorem 4:** Let  $V$  be an  $n$ -dimensional vector space ( $n > 0$ ). Then

- (a) No set of less than  $n$  vectors can span  $V$ ,
- (b) Any set of less than  $n$  linearly independent vectors can be extended to form a basis.
- (c) Any spanning set of more than  $n$  vectors can be pared down to form a basis.

Clearly there are many bases for most vector spaces. However, many common vector spaces have a "standard basis." For example, the standard basis

for  $\mathbb{R}^3$  is  $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right\}$ .

### Homework

1. Determine whether the following vectors are linearly independent in  $\mathbb{R}^2$ .

(a)  $\begin{bmatrix} 3 \\ 8 \end{bmatrix}, \begin{bmatrix} -6 \\ -16 \end{bmatrix}$       (b)  $\begin{bmatrix} -1 \\ 2 \end{bmatrix}, \begin{bmatrix} 5 \\ -2 \end{bmatrix}$       (c)  $\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 4 \\ 4 \end{bmatrix}$

2. Determine whether the following vectors are linearly independent in  $P_3$ .

(a)  $1, x^2, x^2 - 2$       (b)  $1, x, x^2 - 2$       (c)  $x + 2, x + 1, x^2 - 1$

3. Let  $v_1, v_2, \dots, v_k$  be linearly independent vectors in a vector space  $V$ .

- (a) If we add a vector  $v_{k+1}$  to the collection, will we still have a linearly independent set of vectors? Explain.
- (b) If we delete a vector, say  $v_k$ , from the collection, will we still have a linearly independent set of vectors? Explain.

4. Prove that any finite set of vectors that contains the zero vector is linearly dependent.

5. Given  $x_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ ,  $x_2 = \begin{bmatrix} 4 \\ 3 \end{bmatrix}$ , and  $x_3 = \begin{bmatrix} 7 \\ -3 \end{bmatrix}$ . Show that

(a)  $x_1$  and  $x_2$  form a basis of  $\mathbb{R}^2$ .

(b) Why must  $x_1$ ,  $x_2$ , and  $x_3$  be linearly dependent?

(c) What is the dimension of  $\text{Span}(x_1, x_2, x_3)$ ?